



### How this medication works ?

Zepatier® is a combination of two medicines, elbasvir and grazoprevir, indicated for the treatment of hepatitis C virus infection (HCV).

Zepatier® blocks several stages of the virus replication process. This prevents the virus from multiplying in order to promote healing.

### How to take this medication ?

Zepatier® → 1 tablet **once** daily.

This medication must be taken **with food**.

It is important to take Zepatier® regularly, at the same time each day for the duration of the treatment. If you miss doses, it may compromise the success of the treatment and decrease your chances of recovery. Furthermore, the virus responsible for your hepatitis could become resistant and make the medicines less effective or even useless.

The duration of treatment can vary from 8, 12 to 16 weeks depending on :

- Type of virus
- Your liver condition
- Response to previous treatment
- Your kidney condition

### Can I take Zepatier® with my other medicines ?

Tell your healthcare professional especially if you take the following :

- Medicines to prevent transplant rejection like cyclosporine (Neoral)
- Medicines to treat fungal infections like ketoconazole

- Medicines for the treatment of tuberculosis like rifampin (Rifater, Rifadin, Rofact), rifabutin (Mycobutin), or rifapentine
- Medicines to prevent seizures like carbamazepine (Tegretol), oxcarbazepine (Trileptal), phenobarbital (Phenobarb), or phenytoine (Dilantin)
- Medicines to treat pulmonary arterial hypertension like bosentan (Tracleer)
- Medicines to help people stay awake like modafinil (Alertec)
- HIV infection medicines like atazanavir (Reyataz), darunavir (Prezista, Prezcofix), ritonavir (Norvir), lopinavir (Kaletra), tipranavir (Aptivus), efavirenz (Sustiva, Atripla), etravirine (Intelligence), elvitégravir/cobicistat (Stribild, Genvoya)
- Natural product : St. John's Wort (Hypericum perforatum)

These drugs are contraindicated or should be avoided with Zepatier®. Your doctor will suggest an alternative or dose adjustments with closer monitoring.

Other medicines may also interact with Zepatier®, you should tell your doctor about all the health products you take.

Consult your doctor or pharmacist before taking a new prescription medication, over the counter medication, natural product, or even a recreational drug.

### Can I drink alcohol with Zepatier® ?

Ideally, abstinence from alcohol (beer, wine and spirits) is highly recommended for people infected with HCV to protect the liver.

## What are the side effects of this medication ?

It is a drug that is well tolerated in general, the main observed side effects of Zepatier® are mild and do not require stopping treatment.

They include among others :

- Headache
- Fatigue
- Nausea

Tell your doctor and pharmacist about any side effect that bothers you or persists. They can advise you to manage them.

Never stop your treatment without first talking to your doctor or pharmacist.

## Pregnancy and breast feeding

The effects of elbasvir and grazoprevir are not known in pregnant women. Women of childbearing age should use adequate contraception.

Discuss this with your doctor.

The presence of elbasvir and grazoprevir, and their metabolites in breast milk is still unknown. Breastfeeding is therefore not recommended during treatment with Zepatier®.

## Storage

Store Zepatier® tablets in a dry place, at room temperature (15 to 30° C or 59 to 86° F). Avoid damp places such as the bathroom.

Keep this medication in its original container. Keep out of reach and out of sight of children.

## General advice

Be faithful to your appointments with the doctor and the treatment team; they will be able to closely monitor your state of health.


If you have any questions about your medication or need help to schedule when to take them, don't hesitate to ask your doctor or pharmacist who is familiar with what you are taking.

Contact person : \_\_\_\_\_

Telephone : \_\_\_\_\_

@ February 2020, R. Therrien

## I forgot to take a dose !



## Do not double up dose